

Synthesis of Stable 1*H*-Azirines Reinvestigated: A Structural Corrigendum

Klaus Banert,* Manfred Hagedorn, Heiko Peisker


Organic Chemistry, Chemnitz University of Technology, Strasse der Nationen 62, 09111 Chemnitz, Germany
Fax +49(371)53121229; E-mail: klaus.banert@chemie.tu-chemnitz.de

Dedicated to Professor Helmut Quast

Abstract: The isoquinoline-catalyzed synthesis of pretended 1*H*-azirines from phenacyl bromides and *N,N'*-dialkylcarbodiimides was repeated. The products do not possess the structure of antiaromatic 1*H*-azirines, but simple *N*-acyl-*N,N'*-dialkylureas were formed instead. This structural corrigendum was confirmed by the independent synthesis of the known ureas and comparison of their ¹H NMR and ¹³C NMR spectroscopic data in the case of six compounds. Thus, 1*H*-azirines keep their classification as very short-lived intermediates.

Key words: acylation, antiaromatic heterocycles, structure assignment by spectroscopy, ureas

Strained compounds are of special interest because of their increased energy content and the enhanced reactivity, which frequently results from this. For 1*H*-azirines **1** and 2*H*-azirines **2**, it is obvious that both types of heterocycles include considerable ring strain (Scheme 1). However, the properties of **1** and **2** are quite different.¹ A great number of 2*H*-azirines **2**, especially those with R¹ ≠ H, were isolated and characterized by spectroscopic methods in solution or even by X-ray crystallographic structure determination. Although compounds of type **2** are highly reactive, the 2*H*-azirine unit has been found in a few natural products.² On the other hand, only five examples of short-lived 1*H*-azirines **3a,b** were photochemically generated and detected at very low temperatures by IR spectroscopy, which indicated absorptions in the region of 1867–1890 cm⁻¹ attributed to C=C valence vibration.³ Most probably, the push-pull substitution pattern of **3a,b** diminishes the antiaromatic character of the 1*H*-azirine structure and increases the relative stability. Thus, attempts to isolate or observe the parent compound (**1** with R¹ = R² = R³ = H) by cycloaddition⁴ or cycloreversion⁵ approaches and by using argon-matrix isolation technique were unsuccessful and yielded unsubstituted **2** and other isomeric species. Elusive 1*H*-azirine intermediates were merely postulated in several other reactions, which finally led to 2*H*-azirines,⁶ pyrroles,⁷ indoles,⁸ oxazoles,⁹ isoquinolines,¹⁰ ketenimines,¹¹ nitriles,¹² or anilines.¹³ Furthermore, many quantum chemical calculations, that analyzed the energy content,¹⁴ the molecular geometry,¹⁵ the nitrogen inversion barrier,¹⁶ the basicity,¹⁷ and the vibrational frequencies and IR intensities¹⁸ of the parent 1*H*-azirine **1**, have been




Scheme 1 Structures of 1*H*-azirines and 2*H*-azirines

published. All experimental and theoretical results emphasize the properties of the antiaromatic heterocycles **1** as short-lived intermediates, which cannot be isolated at room temperature.

Recently, Alizadeh and Rezvanian reported on the synthesis of 1*H*-azirines **6a–f** by reaction of phenacyl bromides **4a–d** with carbodiimides **5a,b** in the presence of potassium carbonate and a catalytic amount of isoquinoline (10 mol%) in anhydrous acetonitrile (Scheme 2).¹⁹ The products **6a–f** were isolated by chromatography in 65–82% yield as yellow powders and characterized by their sharp melting points (150–176 °C), IR, ¹H NMR, ¹³C NMR and MS data as well as elemental analyses. It was claimed that the antiaromatic heterocycles **6** were formed from **4** and **5** in 35–64% yield even when protic ethanol was used as solvent and strong bases such as potassium hydroxide were utilized. Surprisingly, phenacyl bromide **4e** did not give any product of type **6** after treatment with **5a** or **5b**, and the reaction of **4c** or **4f** with **5a** exclusively led to the simple amides **7a** or **7b**, respectively (yields: 78–80%).¹⁹

We had strong doubts about the antiaromatic structures **6** because the corresponding substances obviously are stable at room temperature, which is absolutely incompatible with 1*H*-azirine moieties.^{1,3–18} The heterocycles **6a–f** should be even less stable than **3a,b** since the electron-withdrawing cyano group of **3** is replaced in **6** by an electron-releasing alkyl group that emphasizes the antiaromatic character of the three-membered ring. Moreover, 1*H*-azirines of type **6** should cause IR signals similar to those of **3a,b**. But IR absorptions within or near to the region of 1867–1890 cm⁻¹ were not found for **6a–f**.



Scheme 2 Synthesis of 1*H*-azirines **6a–f** reported in ref.¹⁹ and the corrected structures **8a–f** of the products from **4** and **5**

Table 1 Comparison of ^{13}C NMR Data^a of the Product from **4b** and **5a** with those of **8c**²⁰

Product from 4b and 5a , ref. ¹⁹	Product from 4b and 5a , this work	8c , ^b this work
20.95	20.94	20.91
22.33	22.32	22.29
42.70	42.69	42.67
50.24	50.18	50.11
55.42	55.42	55.39
113.85	113.81	113.78
122.11	X	X
128.87	128.87	128.84
X	129.17	129.16
133.24	X	X
154.20	154.54	154.51
161.78	161.74	161.71
172.05	172.03	171.93


^aMeasured in CDCl_3 , δ values.

^bPrepared from 4-methoxybenzoic acid and **5a** as described in ref.²¹

When we repeated the reaction of **4a–d** with **5a,b**, we obtained substances with ^1H NMR data which were identical with those reported¹⁹ for **6a–f**. Nearly the same was true for the corresponding ^{13}C NMR spectra. In each of the six cases, however, our ^{13}C

NMR data sets included one quaternary carbon signal less if compared to the ^{13}C NMR spectra published¹⁹ by Alizadeh and Rezvanian. This is shown in Table 1 for the example²⁰ of the product from **4b** and **5a** that obviously cannot have the structure **6c**. Our corrected ^{13}C NMR spectroscopic data excluded the 1*H*-azirines **6a–f** and led to the alternative structures **8a–f**. These *N*-acylureas were described in literature,²¹ and in the cases **8a,c,d** not only ^1H NMR but also ^{13}C NMR spectra were reported.^{21a,b} We prepared **8a–f** for comparison by known procedures and found that ^1H NMR and ^{13}C NMR data were identical with those of the products **8a–f** synthesized from **4a–d** and **5a,b**.²⁰

Scheme 3 Some of the possible reaction mechanisms



to explain the formation of **8**

The intermediates **9**, **14**, and **15** were used to interpret the generation of the alleged 1*H*-azirines **6**, and the ylides **16** were discussed as precursors, which lead to the amides **7a,b** by hydrolysis (Scheme 3).¹⁹ However, these intermediates should alternatively play a role in the formation of **8** from **4** and **5** with loss of one carbon atom. Several reaction mechanisms can be taken into consideration to explain this result. The most simple sequence includes *N*-alkylation of

isoquinoline by **4** and nucleophilic attack of ureas **10**, which are the hydrolysis products of **5**, at the carbonyl group of **9**. This would produce **8** and the salt **11** (X = Br) via the corresponding isoquinolinium ylide. We found that an anhydrous solvent (acetonitrile) is not necessary to prepare **8** from **4** and **5**, and addition of *N,N'*-dicyclohexylurea or one equivalent of water did not prevent the formation of **8**. But no **8d** was observed from **4b** when the co-reactant **5b** was completely exchanged by the corresponding urea. Consequently, carbodiimides **5** are essential to synthesize **8** from **4**.

Cleavage of *N*-phenacylpyridinium salts in aqueous alkaline solution is known for 125 years and can be performed under mild conditions to get benzoic acids and *N*-methylpyridinium salts in good yields.²² This reaction was intensively investigated,²³ utilized in synthesis,²⁴ and also transferred to the corresponding quinolinium^{22a} and isoquinolinium²⁵ salts. Thus, a plausible mechanism to explain the formation of **8** from **4** and **5** includes cleavage of **9** to generate **11** and the benzoates **12**, which can add to **5**. The resulting *O*-acylisoureas **13** are able to undergo a Mumm-like rearrangement to yield **8**. It is well known that treatment of benzoic acids with **5a,b** leads to *N*-acylureas beside the corresponding benzoic anhydrides.^{21c-e,26}

In an alternative reaction mechanism, base-induced transformation of the salts **9** into the stabilized ylides **14** followed by nucleophilic addition at **5** should produce the intermediates **15**. Nucleophilic attack of **10** at the carbonyl group of **15** can yield **8** and the stabilized ylides **18**. On the other hand, intramolecular attack of the NR unit at the carbonyl group of **15** would initiate *N*-acylation and formation of the stabilized ylides **16**. The latter can be hydrolyzed to give **8** and **11**. If **16** adds to a second molecule of **5**, the resulting intermediates **17** should lead to **8** and **18** on hydrolysis.

In summary, we have demonstrated that the reaction of bromides **4** and carbodiimides **5** does not give products with energetic 1*H*-azirine moieties. Consequently, these antiaromatic heterocycles keep their classification as very elusive short-lived intermediates. Instead of alleged 1*H*-azirines **6**, simple *N*-acyl-*N,N'*-dialkylureas **8** were generated from **4** and **5**. The formation of these products is surprising but can be explained by several reaction mechanisms.

Acknowledgements

This is a post-peer-review version from Synlett, 23 (20), K. Banert, M. Hagedorn, H. Peisker, Synthesis of Stable 1*H*-Azirines Reinvestigated: A Structural Corrigendum, 2943-2946, 2012, with permission from Georg Thieme Verlag, self-archived on <http://monarch.qucosa.de> in 2014. The definitive

publisher-authenticated version is available online at: <http://dx.doi.org/10.1055/s-0032-1317494>.

References and notes

- Reviews on 1*H*-azirines and 2*H*-azirines: (a) Backes, J. In *Houben-Weyl, Vol. E16c*; Klamann, D., Ed.; Thieme: Stuttgart, **1992**, 321. (b) Nair, V. In *The Chemistry of Heterocyclic Compounds, Small-Ring Heterocycles, Vol. 42, Part 1*; Hassner, A., Ed.; Wiley: New York, **1983**, 215. (c) Pearson, W. H.; Lian, B. W.; Bergmeier, S. C. In *Comprehensive Heterocyclic Chemistry II, Vol. 1A*; Padwa, A., Ed.; Pergamon: New York, **1996**, 1. (d) Palacios, F.; Ochoa de Retana, A. N.; Martinez de Marigorta, E.; de Los Santos, J. M. *Eur. J. Org. Chem.* **2001**, 2401. (e) Gilchrist, T. L. *Aldrichimica Acta* **2001**, 34, 51. (f) Rai, K. M. L.; Hassner, A. In *Advances in Strained and Interesting Organic Molecules, Vol. 8*; Halton, B., Ed.; Jai Press: Greenwich, **2000**, 187. (g) Fowler, F. W. *Adv. Heterocycl. Chem.* **1971**, 13, 45. (h) Wentrup, C. *Adv. Heterocycl. Chem.* **1981**, 28, 231.
- (a) Miller, T. W.; Tristram, E. W.; Wolf, F. J. *J. Antibiot.* **1971**, 24, 48. (b) Molinski, T. F.; Ireland, C. M. *J. Org. Chem.* **1988**, 53, 2103. (c) Salomon, C. E.; Williams, D. H.; Faulkner, D. J. *J. Nat. Prod.* **1995**, 58, 1463. (d) Stapley, E. O.; Hendlin, D.; Jackson, M.; Miller, A. K.; Hernandez, S.; Mata, J. M. *J. Antibiot.* **1971**, 24, 42. (e) Davis, F. A.; Reddy, G. V.; Liu, H. *J. Am. Chem. Soc.* **1995**, 117, 3651. (f) Davis, F. A.; Liu, H.; Liang, C.-H.; Reddy, G. V.; Zhang, Y.; Fang, T.; Titus, D. D. *J. Org. Chem.* **1999**, 64, 8929.
- Regitz, M.; Arnold, B.; Danion, D.; Schubert, H.; Fusser, G. *Bull. Soc. Chim. Belg.* **1981**, 90, 615.
- (a) Jacox, M. E.; Milligan, D. E. *J. Am. Chem. Soc.* **1963**, 85, 278. (b) Kodama, S. *J. Phys. Chem.* **1988**, 92, 5019.
- Maier, G.; Schmidt, C.; Reisenauer, H. P.; Endlein, E.; Becker, D.; Eckwert, J.; Hess, B. A.; Schaad, L. J. *Chem. Ber.* **1993**, 126, 2337.
- (a) Fowler, F. W.; Hassner, A. *J. Am. Chem. Soc.* **1968**, 90, 2875. (b) Anderson, D. J.; Gilchrist, T. L.; Rees, C. W. *Chem. Commun.* **1969**, 147. (c) Gilchrist, T. L.; Gymer, G. E.; Rees, C. W. *Chem. Commun.* **1971**, 1519. (d) Anderson, D. J.; Gilchrist, T. L.; Gymer, G. E.; Rees, C. W. *J. Chem. Soc., Perkin Trans. 1* **1973**, 550. (e) Gilchrist, T. L.; Gymer, G. E.; Rees, C. W. *J. Chem. Soc., Perkin Trans. 1* **1973**, 555. (f) Atkinson, R. S.; Grimshire, M. J. *J. Chem. Soc., Perkin Trans. 1* **1986**, 1215.
- Gilchrist, T. L.; Gymer, G. E.; Rees, C. W. *J. Chem. Soc., Perkin Trans. 1* **1975**, 1.
- (a) Mitchell, G.; Rees, C. W. *J. Chem. Soc., Chem. Commun.* **1986**, 399. (b) Mitchell, G.; Rees, C. W. *J. Chem. Soc., Perkin Trans. 1* **1987**, 413.
- (a) Huisgen, R.; Blaschke, H. *Chem. Ber.* **1965**, 98, 2985–2997. (b) Meinwald, J.; Aue, D. H. *J. Am. Chem. Soc.* **1966**, 88, 2849.
- Gilchrist, T. L.; Gymer, G. E.; Rees, C. W. *J. Chem. Soc., Chem. Commun.* **1973**, 835.
- Gilchrist, T. L.; Rees, C. W.; Thomas, C. *J. Chem. Soc., Perkin Trans. 1* **1975**, 8.
- (a) Wentrup, C.; Crow, W. D. *Tetrahedron* **1970**, 26, 3965. (b) Thétaz, C.; Wentrup, C. *J. Am. Chem. Soc.* **1976**, 98, 1258.
- Wentrup, C. *Helv. Chim. Acta* **1972**, 55, 1613.
- (a) Clark, D. T. *Theoret. Chim. Acta (Berl.)* **1969**, 15, 225. (b) Hopkinson, A. C.; Lien, M. H.; Yates, K.; Csizmadia, I. G. *Int. J. Quantum Chem.* **1977**, 12, 355.

-
- (c) Bock, H.; Dammel, R. *Chem. Ber.* **1987**, *120*, 1971.
 (d) Alcami, M.; Mó, O.; Yáñez, M. *J. Am. Chem. Soc.* **1993**, *115*, 11074. (e) Lathan, W. A.; Radom, L.; Hariharan, P. C.; Hehre, W. J.; Pople, J. A. *Top. Curr. Chem.* **1973**, *40*, 1.
- (15) (a) Mó, O.; Yáñez, M.; Elguero, J. *J. Mol. Struct. (Theochem)* **1989**, *201*, 17. (b) Byun, Y.-G.; Saebo, S.; Pittman, C. U. *J. Am. Chem. Soc.* **1991**, *113*, 3689. (c) Malar, E. J. P. *Tetrahedron* **1996**, *52*, 4709.
- (16) (a) Budzelaar, P. H. M.; Cremer, D.; Wallasch, M.; Würthwein, E.-U.; Schleyer, P. v. R. *J. Am. Chem. Soc.* **1987**, *109*, 6290. (b) Alcami, M.; de Paz, J. L. G.; Yáñez, M. *J. Comp. Chem.* **1989**, *10*, 468.
- (17) (a) Würthwein, E.-U. *J. Org. Chem.* **1984**, *49*, 2971. (b) Mó, O.; de Paz, J. L. G.; Yáñez, M. *J. Phys. Chem.* **1987**, *91*, 6484.
- (18) Čársky, P.; Hess, B. A.; Schaad, L. J. *J. Am. Chem. Soc.* **1983**, *105*, 396.
- (19) Alizadeh, A.; Rezvanian, A. *Synlett* **2012**, *23*, 859.
- (20) For the complete characterization of **8a–f**, see the Supporting Information.
- (21) (a) Lessene, G.; Smith, B. J.; Gable, R. W.; Baell, J. B. *J. Org. Chem.* **2009**, *74*, 6511. (b) Corriu, R. J. P.; Lanneau, G. F.; Perrot-Petta, M. *Synthesis* **1991**, 954. (c) Neves Filho, R. A. W.; de Oliveira, R. N.; Srivastava, R. M. *J. Braz. Chem. Soc.* **2007**, *18*, 1410. (d) Quintanilla, M. G.; Montero, G.; Hierro, A. G.; Diaz, J. M.; Bassani, S. *Acta Chem. Scand.* **1999**, *53*, 928. (e) Kishikawa, K.; Eida, H.; Kohmoto, S.; Yamamoto, M.; Yamada, K. *Synthesis* **1994**, 173.
- (22) (a) Bamberger, E. *Ber. Dtsch. Chem. Ges.* **1887**, *20*, 3338. (b) Schmidt, E.; van Ark, H. H. *Arch. Pharm.* **1900**, *238*, 321. (c) Babcock, S. H.; Nakamura, F. I.; Fuson, R. C. *J. Am. Chem. Soc.* **1932**, *54*, 4407. (d) Krollpfeiffer, F.; Müller, A. *Ber. Dtsch. Chem. Ges.* **1933**, *66B*, 739. (e) Babcock, S. H.; Fuson, R. C. *J. Am. Chem. Soc.* **1933**, *55*, 2946.
- (23) (a) Kröhnke, F.; Heffé, W. *Ber. Dtsch. Chem. Ges.* **1937**, *70B*, 864. (b) Pearson R. G.; Dillon, R. L. *J. Am. Chem. Soc.* **1948**, *70*, 1933. (c) Pearson, R. G.; Sandy, A. C. *J. Am. Chem. Soc.* **1951**, *73*, 931.
- (24) (a) King, L. C. *J. Am. Chem. Soc.* **1944**, *66*, 894. (b) King, L. C. *J. Am. Chem. Soc.* **1944**, *66*, 1612. (c) King, L. C.; McWhirter, M.; Rowland, R. L. *J. Am. Chem. Soc.* **1948**, *70*, 239. (d) Huffman, J. W.; Zengin, G.; Wu, M.-J.; Lu, J.; Hynd, G.; Bushell, K.; Thompson, A. L. S.; Bushell, S.; Tartal, C.; Hurst, D. P.; Reggio, P. H.; Selley, D. E.; Cassidy, M. P.; Wiley, J. L.; Martin, B. R. *Bioorg. Med. Chem.* **2005**, *13*, 89. (e) Brusotti, G.; Habermann, J.; Azzolina, O.; Collina, S. *Lett. Org. Chem.* **2006**, *3*, 943.
- (25) Kröhnke, F. *Ber. Dtsch. Chem. Ges.* **1933**, *66B*, 604.
- (26) Ślebioda, M. *Tetrahedron* **1995**, *51*, 7829.